

Tris-NTA Biotin

LOT: See product label

EXPIRY DATE: See product label

ORDERING INFORMATION

CAT. NO.	SIZE	PACKAGE CONTENT
BR1001201	100 µg	100 µl Tris-NTA Biotin
BR1001202	1 mg	1 ml Tris-NTA Biotin
BR1001203	1 mg	1 mg Tris-NTA Biotin (powder)

COMPONENT

COMPOSITION

Tris-NTA Biotin	Tris-NTA Biotin (1 mg/ml) in PBS
Tris-NTA Biotin (powder)	Tris-NTA Biotin in powder format, non-dissolved

STORAGE

4°C (until expiry date – see product label)

FEATURES

- A complex of three Ni-NTA groups ensures high-affinity binding of His-tags
- Binding affinity is approximately four orders of magnitude higher than monovalent metal ion chelators
- Protein binding is stoichiometric

APPLICATIONS

- Reversible labeling of proteins or cell surfaces
- Detection and analysis of target molecules
- Immobilization of proteins, lipids and cells on surfaces
- Purification and sample preparation of proteins
- Coupling with microscopic or spectroscopic probes

DESCRIPTION

His-tags are one of the most commonly used tags for protein expression analysis. Conventional metal ion chelators, such as nitrilotriacetic acid (NTA) and iminodiacetic acid (IDA), bind His-tags with low affinities in the range of 10 µM. The biotechrabbit Tris-NTA complexes three NTA groups that together bind a 6×His-tag with an affinity that is four orders of magnitude higher (1 nM) than is possible with conventional chelators (10 µM). The binding of His-tags is stoichiometric and single-molecule detection is possible. Binding is reversible: bound His-tags can be released with imidazole or ethylenediaminetetraacetic acid (EDTA).


biotechrabbit™ Tris-NTA is available with a free amine group or conjugated to biotin. It can be used in a large range of applications, including protein detection and labeling, coupling proteins, lipids or cells to surfaces, protein purification and reversible protein modification.

Tris-NTA Biotin

SPECIFICATIONS

CHEMICAL NAME	Biotin Tris-NTA (trifluoroacetic acid salt)
STABILITY	24 months
STORAGE CONDITIONS	Store at 4°C
METHOD FOR DETERMINING IDENTITY	Mass spec analysis
METHOD FOR DETERMINING PURITY	HPLC
CAS NUMBER	1070867-85-4
MOLECULAR FORMULAR	C ₅₉ H ₉₃ N ₁₁ O ₂₅ S
MOLECULAR WEIGHT	1388.49
SOURCE	Synthetic
PURITY	>95% (HPLC)
FORM	1mg/ml solution in PBS or powder

CHEMICAL STRUCTURE



APPLICATION EXAMPLES

High-Affinity Adaptors for Switchable Recognition of Histidine-Tagged Proteins.

Lata et al., *J. Am. Chem. Soc.*, 2005, 127, 10205–10215

Specific and Stable Fluorescence Labeling of Histidine-Tagged Proteins for Dissecting Multi-Protein Complex Formation.

Lata et al., *J. Am. Chem. Soc.*, 2006, 128, 2365–2372

Noncovalent, Site-Specific Biotinylation of Histidine-Tagged Proteins.

Reichel et al., *Anal Chem.*, 2007, 79, 8590–600

Identifying Modulators of Protein-Protein Interactions Using Photonic Crystal Biosensors.

Heeres et al., *J Am Chem Soc.* 2009, 131:18202–18203

Tris-Nitrilotriacetic Acids of Sub-nanomolar Affinity Toward Hexahistidine Tagged Molecules.

Huang et al., *Bioconjug Chem.*, 2009, 20: 1667–1672

Four-color single-molecule fluorescence with noncovalent dye labeling to monitor dynamic multimolecular complexes.

DeRocco et al., *BioTechniques* 2010, 49: 807-816

In situ assembly of macromolecular complexes triggered by light.

Grunwald et al., *PNAS*, 2010, 107: 6146-6151

Quantum-Yield-Optimized Fluorophores for Site-Specific Labeling and Super-Resolution Imaging.

Grunwald, et al., *J. Am. Chem. Soc.*, 2011, 133, 8090–8093.

Co- and distinct existence of Tris-NTA and biotin functionalities on individual and adjacent micropatterned surfaces generated by photo-destruction.

Biswas et al., *Soft Matter*, 2014, 10, 2341–2345

High-affinity gold nanoparticle pin to label and localize histidine tagged protein in macromolecular assemblies.

Anthony et al., *Structure*, 2014, 22: 628–635

Tris-NTA Biotin

CERTIFICATE OF ANALYSIS

Quality Control

Identity

Identity of the substance was determined by MS analysis. The identity was consistent with the reference substance.

Purity

The purity of Tris-NTA Biotin was determined by HPLC. Purity was >95%.

Quality confirmed by: Head of Quality Control

SAFETY INSTRUCTIONS

For safety instructions please see Safety Data Sheets (SDS)/Sicherheitshinweise finden Sie in den SDS unter: <http://www.biotechrabbit.com/support/documentation.html>.

USEFUL HINTS

- Visit Applications at www.biotechrabbit.com for more products and product selection guides.
- Most biotechrabbit products are available in custom formulations and bulk amounts.

CONTACT BIOTECHRABBIT

biotechrabbit GmbH
Volmerstr. 9a
12489 Berlin, Germany

info@biotechrabbit.com
support@biotechrabbit.com
www.biotechrabbit.com

Phone: +49 30 555 7821-10
Fax: +49 30 555 7821-99



Legal Disclaimer and Product Use Limitation

Purchase of product does not include a license to perform any patented applications; therefore it is the sole responsibility of users to determine whether they may be required to engage a license agreement depending upon the particular application in which the product is used. This product was developed, manufactured, and sold for in vitro use only. It is not suitable for administration to humans or animals.

The use of this product for research purposes is covered by a license from QIAGEN Group. No rights to use this product to perform or offer diagnostic, companion diagnostic, commercial testing or other commercial services for money or money's worth are granted by the supply of this product expressly, by implication or by estoppel. Should you wish to use this product for any other purpose not covered by this license, please contact QIAGEN Corporate Business Development at bd@qiagen.com.

Trademarks: biotechrabbit™ (biotechrabbit GmbH, QIAGEN® (QIAGEN Group).

valid from 24.08.2016