


α2,3 Sialidase contents

Catalog #	Description	Size	M. W.	Purity	pH	Storage
GE0302	α2,3 Sialidase	5,000 units, lyophilized	77,512	> 95%	7.0-7.5 optimal	-20°C, up to 12 months
BA0801	10X Reaction Buffer 4	1 mL			7.5	4 to 25°C

This product is for research use only and not for resale or for any use in the manufacture of a therapeutic or for any diagnostic purpose.

Product description: This product is a recombinant neuraminidase (exo-α-sialidase; glycosyl hydrolase family GH33, EC #3.2.1.18), cloned from *Streptococcus pneumoniae* and expressed in *Escherichia coli* with an N-terminal 8xHis tag. The 8xHis tag may be removed by digestion with FasTEV[™] (Cat #GE0501), a TEV protease with enhanced stability and catalytic activity.

This product preferentially releases terminal α2,3-linked N-acetylneuraminic acid (Neu5Ac) from oligosaccharides, complex carbohydrates, and glycoproteins.



This product does not contain any detectable activities or proteases or other glycosidases.

Unit definition: One unit is defined as the amount of α2,3 Sialidase required to catalyze the release of 1 nanomole of p-nitrophenol (pNP) from 2-O-(p-nitrophenyl)-α-D-N-acetylneuraminic acid (pNP-Neu5Ac) in 1 min at 37°C in 100 μL 1X Reaction Buffer 4 (50 mM Tris-HCl, 100 mM NaCl, pH 7.5).

Product reconstitution: Dissolve the lyophilized product in 100 μL of molecular grade water to make a 50,000 units/ml (Cat #GE0302) solution in enzyme storage buffer (50 mM EPPS, 100 mM NaCl, pH 7.5). Once reconstituted, store at 4°C for up to 5 days or -20°C for up to 6 months. Aliquoting is recommended to avoid repeated freeze-thaw cycles.

Suggested protocol for protein desialylation:

- Mix the following components in a microfuge tube:

Glycoprotein (e.g., fetuin; user supplied)	1 nanomole (2-100 μg)
10X Reaction Buffer 4 (Cat #BA0801)	10 μL
α2,3 Sialidase (Cat #GE0302)	1.0 μL (50 units)
Molecular grade water	to 100 μL final volume
- Incubate at 37°C for 1 h.
- Analyze by Western blot or other method to determine the extent of desialylation on the substrate. Suggested 1° probes for Western blot analysis: biotinylated SiaFind[™] α2,3-Specific Lectenz[®] (Cat #SK2301B), SiaFind[™] Pan-Specific Lectenz[®] (Cat #SK0501B), or SiaFind[™] Pan-Specific Lectenz[®] 2.0 (Cat #SK0502B).

Reference: Gut H, et al. FEBS Lett. 2008 Oct 15;582(23-24):3348-52. PMID: 18775704.

Note: Reactions may be scaled up to accommodate larger amount and volume of substrate. Amount of enzyme and reaction time may vary for different substrates. Titration of the amount of enzyme in a reaction is recommended for each new substrate.