



Certificate of Analysis

www.tocris.com

Product Name: LY 215840 Catalog No.: 1523 Batch No.: 1

CAS Number: 137328-52-0

IUPAC Name: [8β(1S,2R)]-N-(2-Hydroxycyclopentyl)-6-methyl-1-(1-methylethyl)-ergoline-8-carboxamide

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{24}H_{33}N_3O_2$. ¹/₄ H_2O

Batch Molecular Weight: 400.04
Physical Appearance: White solid

Solubility: DMSO to 100 mM ethanol to 25 mM

Storage: Store at +4°C

Batch Molecular Structure:

2. ANALYTICAL DATA

HPLC: Shows 98.9% purity

¹H NMR: Consistent with structure Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 72.06 8.44 10.5 Found 72.02 8.34 10.44



Product Information

Print Date: Jan 16th 2012

www.tocris.com

Product Name: LY 215840 Catalog No.: 1523 Batch No.: 1

CAS Number: 137328-52-0

IUPAC Name: [8β(1S,2R)]-N-(2-Hydroxycyclopentyl)-6-methyl-1-(1-methylethyl)-ergoline-8-carboxamide

Description:

 $5\text{-HT}_2/5\text{-HT}_7$ receptor antagonist. Reduces stimulatory effect of serotonin on aldosterone secretion; also reduces 5-HT-induced cAMP formation (pK_R = 8.26).

Physical and Chemical Properties:

Batch Molecular Formula: C₂₄H₃₃N₃O₂.¼H₂O

Batch Molecular Weight: 400.04 Physical Appearance: White solid

Minimum Purity: >98%

Batch Molecular Structure:

Storage: Store at +4°C

Solubility & Useage Info:

DMSO to 100 mM ethanol to 25 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Cushing *et al* (1996) LY215840, a high-affinity 5-HT7 receptor ligand, blocks serotonin-induced relaxation in canine coronary artery. J.Pharmacol.Exp.Ther. **277** 1560. PMID: 8667223.

Lenglet *et al* (2002) Activation of 5-HT₇ receptor in rat glomerulosa cells is associated with an increase in adenylyl cyclase activity and calcium influx through T-type calcium channels. Endocrinology *143* 1748. PMID: 11956157.

Louiset *et al* (2006) Expression of serotonin7 receptor and coupling of ectopic receptors to protein kinase A and ionic currents in adrenocorticotropin-independent macronodular adrenal hyperplasia causing Cushing's syndrome. J.Clin.Endocrinol.Metab. *91* 4578. PMID: 16954157.

