

DESCRIPTION

Species Reactivity	Human
Specificity	Detects Recombinant Human IL-18 R α /IL-1 R5 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) IL-1 R1, rhIL-1 RII, rhIL-1 R3, rhIL-1 R4, rhIL-1 R6, rhIL-1 R7, rhIL-1 R8, rhIL-1 R9, rhSIGIRR or recombinant mouse IL-18 R α is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 70625
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human IL-18 R α /IL-1 R5 Glu20-Arg329 Accession # Q13478
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 μ g/10 ⁶ cells	Human peripheral blood mononuclear cells treated with PHA and Recombinant Human IL-2 (Catalog # 202-IL)

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Interleukin 18 (IL-18) is a member of the IL-1 family of cytokines and shares numerous immunoregulatory functions with IL-12. The functional IL-18 receptor complex is composed of two subunits designated IL-18 R α (also termed IL-1 R5 and IL-1 Rrp) and IL-18 R β (also termed IL-1 R7 and AcPL). Both IL-18 R α and IL-18 R β belong to the IL-1 receptor superfamily. Although IL-18 R by itself binds IL-18 with low affinity and IL-18 R β does not bind IL-18 *in vitro*, co-expression of IL-18 R α and IL-18 R β is required for high affinity binding and IL-18 responsiveness. Human IL-18 R cDNA encodes a 541 amino acid (aa) precursor type I membrane protein with a hydrophobic signal, an extracellular domain comprised of three immunoglobulin-like domains, a transmembrane domain and a cytoplasmic region of approximately 200 aa. Human and mouse IL-18 R share 65% amino acid sequence homology. IL-18 R is widely expressed in numerous tissues including spleen, thymus, leukocyte, liver, lung, heart, small and large intestine, prostate and placenta.

References:

1. Parnet, P. *et al.* (1996) J. Biol. Chem. **271**:3967.
2. Torigoe, K. *et al.* (1997) J. Biol. Chem. **272**:25737.
3. Born, T.L. *et al.* (1998) J. Biol. Chem. **273**:29445.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.