





# Dolichos Biflorus Agglutinin (DBA), Unconjugated

## Product Images




## Short Description

*Dolichos biflorus* agglutinin has a carbohydrate specificity toward  $\alpha$ -linked *N*-acetylgalactosamine. It has been used to establish secretor status in blood group A individuals by hemagglutination inhibition techniques and for blood typing. This lectin has also been used as a general marker of developing renal collecting ducts.

## Additional Information

Unit Size	5 mg
Applications	Immunohistochemistry / Immunocytochemistry, Immunofluorescence, Blotting Applications, Glycobiology
Recommended Usage	Although many buffers can be employed for reconstituting and diluting this lectin, 10 mM HEPES buffered saline, pH 8.5, 0.1 mM $\text{CaCl}_2$ is recommended. For preserving solutions stored at 4 °C, 0.08% sodium azide can be used. DO NOT VORTEX!
Recommended Storage	2-8 °C
Conjugate	Unconjugated
Sugar Specificity	N-Acetylgalactosamine

## Products in this set



### Dolichos Biflorus Agglutinin (DBA), Unconjugated

This lectin has a carbohydrate specificity toward  $\alpha$ -linked *N*-acetylgalactosamine. It has been used to establish secretor status in blood group A individuals by hemagglutination inhibition techniques and for blood typing. This lectin has also been used as a general marker of developing renal collecting ducts.

