



Common γ_c -Chain Family Cytokines Regulate Immune System Functions



IL-7 IL-7 Receptor-expressing cells:
T cells, pre-B cells, dendritic cells



IL-4 IL-4 Receptor-expressing cells:
T cells, B cells, NK cells, basophils, mast cells



IL-2 IL-2 Receptor-expressing cells:
T cells, B cells, NK cells, monocytes, macrophages



γ_c Family Cytokines Have Unique & Overlapping Effects on Different Immune Cell Types


Cytokines belonging to the common cytokine receptor γ_c family include IL-2, IL-4, IL-7, IL-9, IL-15, and IL-21. Members of this family signal through receptor complexes that contain the γ_c /IL-2 R β subunit. The γ_c subunit associates with different cytokine-specific receptor subunits to form unique heterodimeric receptors for IL-4, IL-7, IL-9, and IL-21, or associates with both IL-2/IL-15 R β and IL-2 R α or IL-15 R α to form heterotrimeric receptors for IL-2 or IL-15, respectively. γ_c family cytokines generally activate three major signaling pathways that promote cellular survival and proliferation, the PI 3-K-Akt pathway, the RAS-MAPK pathway, and the JAK-STAT pathway. Differences in the expression patterns of the cytokines or their unique receptor components, along with the activation of different STAT proteins may account for some of the distinct effects mediated by γ_c family cytokines.

Signaling by γ_c family cytokines plays a major role in regulating the development, survival, proliferation, differentiation and/or function of cells of the immune system. The importance of the γ_c family cytokines for the establishment and maintenance of the immune system is


emphasized by the fact that mutations in γ_c /IL-2 R β in humans are associated with a disease known as X-linked severe combined immunodeficiency (XSCID), which is characterized by the absence of T cells and natural killer (NK) cells, and the presence of non-functional B cells. Knockout studies in mice have demonstrated that the lack of T cell and NK cell development in this disease can be primarily attributed to the respective loss of IL-7 and IL-15 signaling, while the loss of both IL-4 and IL-21 signaling leads to defective B cell function. Similar studies revealed that in contrast to humans, B cell development in mice also requires IL-7 signaling. Several additional unique and overlapping effects of the γ_c family cytokines on different immune cell types have been documented. A number of these effects are highlighted here to demonstrate the central role that γ_c family cytokines play in controlling immune system functions. Understanding the mechanisms by which these cytokines act and how their signaling pathways can be regulated may have therapeutic implications not only for a variety of immunodeficient disease states, but also for disorders resulting from aberrant or exaggerated immune system activation.

γ_c FAMILY CYTOKINE-INDUCED EFFECTS ON:


T Cells B Cell NK or NKT Cells Mast Cells or Basophils Dendritic Cells Epithelial Cells



IL-15 IL-15 Receptor-expressing cells:
T cells, NK cells, NKT cells, dendritic cells, monocytes



IL-21 IL-21 Receptor-expressing cells:
T cells, B cells, NK cells, NKT cells, dendritic cells, lymphoid tissues



This illustration represents general processes suggested in the scientific literature and is not to be considered comprehensive nor definitive. ©2011, 2015 R&D Systems