


Catalog Number : KRN7000

Lot Number : 5E-12A

Size : 1mg

Chemical structure



Chemical name

(2S,3S,4R)-1-O-(α D-galactosyl)-N-hexacosanoyl-2-amino-1,3,4-octadecanetriol

Molecular formula C₅₀H₉₉NO₉

Molecular weight 858.34

Purity >99% (TLC)

Proton NMR Pass

Biological Activity Pass

Appearance White to off-white powder.

Solubility KRN7000 is practically insoluble in water, methanol or ethanol, very slightly soluble in tetrahydrofuran, slightly soluble in pyridine, and practically insoluble in other organic solvents.

Example of how to dissolve for biological assay

KRN7000 can be dissolved as follows for various biological examinations:

In vivo administration : KRN7000 can be dissolved in a vehicle described by Giaccone et al.(10). Such a vehicle consists of 5.6% sucrose, 0.75% L-histidine, and 0.5% Tween 20. After mixing, the solution should be heated at 80°C until the material becomes completely dissolved and disappeared. This solution can be lyophilized and lyophilized powder can be reconstituted easily by pure water.

In vitro use : KRN7000 can be dissolved in DMSO at a concentration of 1 mg/mL. The solution should be heated at 80°C to get a clear solution.

Storage Short term storage +4°C, Long term storage -20°C

Shipping **Shipped on Blue ice**

License This product is licensed by Kyowa Hakko Kirin Co., Ltd.

Warning Research use only. Not for use in humans.

Background

α Galactosylceramide(α Gal-Cer;KRN7000), an agelasphin derivative developed by Kirin Brewery Co., Ltd., is a biological response modifier (BRM). Agelasphins was isolated from an extract

of the marine sponge, *Agelas mauritianus*, as active substances. They are compounds with α -Galactosylceramide structures, that is, galactose combined with ceramide in an α configuration.

α Gal-Cer; KRN7000, a chemically synthesized α Galactosylceramide, is a specific ligand for human and mouse natural killer T (NKT) cells, KRN7000 exhibits potent antitumor activity in various kinds of *in vivo* murine experimental models including subcutaneously implanted model and metastatic models in the liver and lung. In the liver metastatic models, treatment with KRN7000 suppressed the growth of tumors and prolonged the survival term of tumor-bearing mice. KRN7000 has been reported to show various immunological effects in infectious disease, autoimmune disease, and graft versus host disease in mice.

References

- 1) Natori, T., et al. Agelasphins, Novel Antitumor and Immunostimulatory Cerebrosides from the Marine Sponge *Agelasmauritianus*, *Tetrahedron Lett.*, 34, 5591-5592, (1993).
- 2) Akimoto, K., et al. Synthesis and Stereochemistry of Agelasphin-9b, *Tetrahedron Lett.*, 34, 5593-5596, (1993).
- 3) Natori, T., et al. Agelasphins, novel antitumor and immunostimulatory cerebrosides from the marine sponge *Agelas mauritianus* *Tetrahedron*, 50, 2771-2784, (1994).
- 4) Motoki, K., et al. Immunostimulatory and Antitumor Activities of Monoglycosylceramides Having Various Sugar Moieties *Biol. Pharm. Bull.*, 18, 1487-1491, (1995).
- 5) Morita, M., et al. Syntheses of α -, β -monoglycosylceramides and four diastereomers of an α -galactosylceramide *Bioorganic Med. Chem. Lett.*, 5, Pages 699-704
- 6) Motoki, K., et al. Antitumor activities of α -, β -monogalactosylceramides and four diastereomers of an α -galactosylceramide *Bioorganic Med. Chem. Lett.*, 5, 705-710, (1995).
- 7) Kawano, T., et al. CD1d-restricted and TCR-mediated activation of α 14 NKT cells by glycosylceramides. *Science*, 278, 1626-1629, (1997).
- 8) Kronenberg, M. Toward an understanding of NKT cell biology: progress and paradoxes. *Annu Rev Immunol.*, 23 :877-900 (2005).
- 9) Yamaguchi, Y., et al. Enhancing effects of (2S,3S,4R)- 1-O- (α -D-galactopyranosyl)-2-(N-hexacosanoylamino) -1,3,4-octadecanetriol (KRN7000) on antigen-presenting function of antigen-presenting cells and antimetastatic activity of KRN7000-pretreated antigen-presenting cells. *Oncology Res.*, 8, 399-407, (1996).
- 10) Giaccone, et al. A Phase I Study of the Natural Killer T-Cell Ligand α -Galactosylceramide (KRN7000) in Patients with Solid Tumors *Clinical Cancer Res.* 8:3702–3709, (2002)
- 11) Shimizu, K., et al. Cross-presentation of glycolipid from tumor cells loaded with α -galactosylceramide leads to potent and long-lived T cell mediated immunity via dendritic cells. *S, J Exp Med.*, 204 : 2641-2653 (2007)
- 12) Ishii Y., et al. α -galactosylceramide-driven immunotherapy for allergy. *Front Biosci.* 2008 May 1;13:6214-28.
- 13) Kamijuku H. et al. Mechanism of NKT cell activation by intranasal coadministration of α -galactosylceramide, which can induce cross-protection against influenza viruses, *Mucosal Immunol.* 2008 May;1(3):208-218.